


## Clinical evidence references



## Clinical evidence references

Throughout the entire care pathway, resident/patient mobility is one of the most pivotal constants that impacts decision making. Assessment and communication tools for patient/resident mobility can make therefore all the difference when providing safe and high-quality care for patients and caregivers alike. They must be easy-to-use, validated, and evidence-based.

This reference list details the clinical and research evidence base supporting both Arjo's Positive Eight, the link between mobility and quality of work and care, and the Mobility Gallery with its close association to the validated functional mobility assessment tools and International guidelines.











D

References

- Achterberg WP, van Campen C, Pot AM, Kerkstra A, Ribbe MW. Effects of the Resident Assessment Instrument on the care process and health outcomes in nursing homes. A review of the literature. Scandinavian journal of rehabilitation medicine. 1999 Sep;31(3):131-7.
- 2. Achterberg WP, Holtkamp CC, Kerkstra A, Pot AM, Ooms ME, Ribbe MW. Improvements in the quality of co-ordination of nursing care following implementation of the Resident Assessment Instrument in Dutch nursing homes. Journal of Advanced Nursing. 2001 Jul 22;35(2):268-75.
- 3. Centers for Medicare & Medicaid Services.
  Long-Term Care Facility Resident Assessment
  Instrument 3.0 User's Manual. Version 1.16.
  Resident Assessment Instrument (RAI) 2018
  Oct. [cited 10 September 2020]. Available from:
  https://downloads.cms.gov/files/1-MDS-30RAI-Manual-v1-16-October-1-2018.pdf
- 4. Dijkstra A et al. Measuring care dependency with the care dependency scale. A manual. Eurecare. Guidebook for architects and planners. Arjo. 2006.

- Doroszkiewicz H, Sierakowska M, Muszalik M. Utility of the Care Dependency Scale in predicting care needs and health risks of elderly patients admitted to a geriatric unit: a crosssectional study of 200 consecutive patients. Clinical Interventions in Aging. 2018;13:887-94.
- 6. González-Seguel F, Corner EJ, Merino-Osorio C. International classification of functioning, disability, and health domains of 60 physical functioning measurement instruments used during the adult intensive care unit stay: A scoping review. Physical therapy. 2019 May 1;99(5):627-40.
- 7. Hignett S, Fray M, Battevi N, Occhipinti E, Menoni O, Tamminen-Peter L, Waaijer E, Knibbe H, Jäger M. International consensus on manual handling of people in the healthcare sector: Technical report ISO/TR 12296. International Journal of Industrial Ergonomics. 2014 Jan 1;44(1):191-95.
- 8. Hoben M, Poss JW, Norton PG, Estabrooks CA. Oral/dental items in the resident assessment instrument minimum data set 2.0 lack validity: results of a retrospective, longitudinal validation study. Population health metrics. 2016 Dec 1;14(1):36.



- Hutchinson AM, Milke DL, Maisey S, Johnson C, Squires JE, Teare G, Estabrooks CA. The resident assessment instrument - minimum data set 2.0 quality indicators: a systematic review. BMC health services research. 2010 Dec 1;10(1):166.
- ISO/TR 12296:2012 Ergonomics Manual handling of people in the healthcare sector [Internet]. ISO. 2012 [cited 10 September 2020]. Available from: https://www.iso.org/ standard/51310.html.
- Lahmann NA, Tannen A, Kuntz S, Raeder K, Schmitz G, Dassen T, Kottner J. Mobility is the key! Trends and associations of common care problems in German long-term care facilities from 2008 to 2012. International journal of nursing studies. 2015 Jan 1;52(1):167-74.
- Low LF, Fletcher J, Goodenough B, Jeon YH, Etherton-Beer C, MacAndrew M, Beattie E. A systematic review of interventions to change staff care practices in order to improve resident outcomes in nursing homes. PloS one. 2015 Nov 11;10(11):e0140711.
- 13. Pieber K, Herceg M, Paternostro-Sluga T, Pablik E, Quittan M, Nicolakis P, Fialka-Moser V, Crevenna R. Reliability, validity, sensitivity and internal consistency of the ICF based Basic Mobility Scale for measuring the mobility of patients with musculoskeletal problems in the acute hospital setting: a prospective study. BMC musculoskeletal disorders. 2015 Dec;16(1):1-8.
- Poss JW, Jutan NM, Hirdes JP, Fries BE, Morris JN, Teare GF, Reidel K. A review of evidence on the reliability and validity of Minimum Data Set data. Healthcare Management Forum. 2008 Apr (Vol. 21:1):133-39.

- 15. Prodinger B, Stucki G, Coenen M, Tennant A. The measurement of functioning using the International Classification of Functioning, Disability and Health: comparing qualifier ratings with existing health status instruments. Disability and rehabilitation. 2017 Oct 8:1-8.
- Resident Assessment Instrument (RAI). RAI-MDS 2.0 [Internet]. Interiorhealth.ca. 2020 [cited 10 September 2020]. Available from: https://www.interiorhealth.ca/sites/Partners/ SeniorsCare/RAI/Pages/default.aspx.
- 17. Stucki G. International Classification of Functioning, Disability, and Health (ICF): a promising framework and classification for rehabilitation medicine. American journal of physical medicine & rehabilitation. 2005 Oct 1;84(10):733-40.
- 18. Wald HL, Ramaswamy R, Perskin MH, Roberts L, Bogaisky M, Suen W, Mikhailovich A. The case for mobility assessment in hospitalized older adults: American Geriatrics Society white paper executive summary. Quality and Performance Measurement Committee of the American Geriatrics Society. Journal of the American Geriatrics Society. 2018 Jan:11-6.
- Waters TR, Putz-Anderson V, Garg A.
   National Technical Implementation Service.
   1991, Scientific Support Documentation for the Revised /99/ NJOSH Lifting Equation, PB91-226274 (US Department of Commerce, Springfield, VA).
- 20. World Health Organization (WHO). International Classification of Functioning, Disability, and Health: ICF-2. Geneva: World Health Organization; 2001.





At Arjo, we are committed to improving the everyday lives of people affected by reduced mobility and age-related health challenges. With products and solutions that ensure ergonomic patient handling, personal hygiene, disinfection, diagnostics, and the effective prevention of pressure injuries and venous thromboembolism, we help professionals across care environments to continually raise the standard of safe and dignified care. Everything we do, we do with people in mind.

www.arjo.com

